

RESEARCH

Open Access



Public opinions about supervised smoking facilities for crack cocaine and other stimulants

Carol Strike^{1*}, Nooshin Khobzi Rotondi², Tara Marie Watson¹, Gillian Kolla¹ and Ahmed M. Bayoumi^{3,4,5,6}

Abstract

Background: The purpose of this study was to estimate awareness and opinions about supervised smoking facilities (SSFs) for smoking crack cocaine and other stimulants and make comparisons with awareness and opinions about supervised injection facilities (SIFs) in Ontario, Canada.

Methods: We used data from a 2009 telephone survey of a representative adult sample. The survey asked about awareness of, and level of support for, the implementation of SSFs and SIFs. Data were analysed using statistical models for complex survey data, which account for stratified sampling and incorporate sampling weights.

Results: A total of 1035 participated in the survey. Significantly fewer had knowledge about SSFs (17.9 %) than about SIFs (57.6 %). Fewer strongly agreed with implementation of SSFs (19.6 %) than SIFs (28.3 %). Just over half (51.1 %) of participants somewhat agreed or disagreed, 15.7 % strongly agreed, and 10.6 % strongly disagreed with implementing both SSFs and SIFs.

Conclusions: Members of the public in Ontario had little knowledge of SSFs compared to SIFs. Recent federal government changes in Canada may provide the leadership environment necessary to ensure that innovative, evidence-based harm reduction programs such as SSFs are developed and implemented.

Keywords: Supervised smoking facilities, Public opinion, Telephone survey, General population, Adult, Canada

Background

The past thirty years have seen an expansion in public health programming to prevent drug-related harms for people who inject illicit drugs such as heroin, other opioids, cocaine and crack cocaine, and crystal methamphetamine. Well-known examples include needle and syringe programs and methadone maintenance treatment, both of which are effective at reducing human immunodeficiency virus (HIV) transmission [1, 2]. In recent years, there have been increasing calls to base drug policy and programming on evidence instead of ideology [3], including for example the implementation of supervised consumption facilities. These programs have been implemented in some countries to provide a

hygienic environment where clients can inject and/or smoke illicitly obtained drugs [4].

Supervised injection facilities (SIFs) are a common type of consumption facility and a programmatic innovation with the goal of addressing injection-related harms and risks, including: reducing transmission of HIV, hepatitis C virus (HCV), and other blood-borne infections; decreasing morbidity, overdose and mortality associated with public drug use; minimising public disturbances and drug-related litter; and increasing contact between people who use drugs and health and social services [5–7]. Across the world, there are an estimated 90 supervised consumption facilities implemented where problem drug use continues despite the availability of drug treatment, needle and syringe programs, social services, and attempts by police to reduce drug-related public disorder and drug markets [8, 9]. A large and growing literature from Australia, Europe, and Canada has demonstrated numerous benefits of SIFs [10–16].

* Correspondence: carol.strike@utoronto.ca

¹Dalla Lana School of Public Health, University of Toronto, 155 College Street, Toronto, Canada

Full list of author information is available at the end of the article

Across North America and Europe, smoking of stimulants such as crack cocaine is prevalent and associated with many negative health and social effects [17–21]. Reports suggest that the sharing of smoking equipment is common [22] and potentially exposes smokers to HCV and tuberculosis [23–26]. However, public health responses to these harms have been slower to develop than those related to injection drug use [22]. Nevertheless, some Canadian public health departments distribute low-cost or free safer crack cocaine smoking equipment (e.g., glass stems and mouthpieces) to reduce equipment sharing and potential injuries to the hands and mouth from using damaged smoking equipment [27]. In the United Kingdom, where smoking of heroin is more prevalent, some programs now distribute foil sheets designed to encourage the transition from injecting to heroin smoking [28]. Similar to SIFs, supervised smoking facilities (SSFs) operate in some European countries to reduce illicit drug smoking-related harms such as disease transmission, morbidity associated with public drug use, public disorder, drug-related litter, and limited access to health and social services [6].

There are no legally sanctioned SSFs in North America, although there have been calls for implementation of these programs [29, 30]. Results from an ethnographic study of an unsanctioned SSF in Vancouver, Canada operated by a drug user-led organisation showed the potential of SSFs to attract highly vulnerable people who smoke crack cocaine and to minimise client exposure to interpersonal violence [31]. However, the study showed only modest potential to reduce crack pipe sharing [31]. In other cities in Canada where crack cocaine smoking is prevalent, over two-thirds of people who smoke crack cocaine reported a willingness to use a SSF [5, 30].

Interest in these facilities has been raised by both public health officials and advocates in Canadian cities where crack cocaine smoking is prevalent [5, 32]. To our knowledge, there are no published studies that report public perception of SSFs or reports comparing opinions about SSFs relative to SIFs. However, public opinion is a factor in decision making regarding implementation of public health programs [33, 34]. The goal of our study was to estimate the level of public awareness of SSFs in general and relative to SIFs. In addition, we estimated the level of public agreement or lack thereof regarding implementation of SSFs, both in general and with respect to specific SSF goals.

Methods

Study design and data source

We analysed data from the Centre for Addiction and Mental Health (CAMH) Monitor survey, an annual cross-sectional survey that used computer-assisted telephone interviews to ask participants from Ontario, Canada about

drug issues and policies as well as substance use and mental health. We used the 2009 survey, in which about 2000 adults were selected. Sampling followed a two-stage probability design. First, households were sampled using randomly selected telephone numbers within six regional strata. Second, the person within each household who was both age 18 or over and had the most recent birthday was selected. Fluency in English or French was required for eligibility. All participants provided informed consent. Additional information is available in the CAMH Monitor technical guide [35]. This study was approved by the Research Ethics Board at the Centre for Addiction and Mental Health.

For six of the months in 2009, survey participants were first asked about their opinions regarding SIFs and next asked about their opinions related to SSFs [36]. Prior to being asked for their opinions, participants were told that they would be asked questions about SIFs and were provided with the following description: “The Vancouver supervised injection facility, ‘Insite,’ provides a place supervised by health care workers for drug users to inject their drugs. Several other cities in Canada are considering starting up similar programs.” Participants were asked if they strongly agreed, somewhat agreed, somewhat disagreed or strongly disagreed with each of the following statements:

1. Supervised injection facilities should be made available to injection drug users, to encourage safer drug injection.
2. Supervised injection facilities should be made available if it can be shown that they reduce overdose deaths or infectious disease among users.
3. Supervised injection facilities should be made available if they can increase drug users’ contact with health and social workers.
4. Supervised injection facilities should be made available if it can be shown that they reduce neighbourhood problems related to injection drug use.

After asking about SIFs, interviewers informed participants that, “Several cities in Canada are considering starting up similar facilities supervised by health care workers for drug users to smoke drugs like crack cocaine and methamphetamine. The next few questions are about your views on these facilities.” Specifically, participants were asked:

1. Have you ever read, seen or heard any information about supervised smoking facilities?
2. Please indicate if you strongly agree, somewhat agree, somewhat disagree or strongly disagree with the following statements:
 - a. Supervised smoking facilities should be made available to people who smoke drugs like crack

- cocaine and methamphetamine to encourage safer drug consumption
- b. Supervised smoking facilities should be made available if it can be shown that they reduce infectious disease among to people who smoke drugs like crack cocaine and methamphetamine
 - c. Supervised smoking facilities should be made available if they can increase drug users' contact with health and social workers
 - d. Supervised smoking facilities should be made available if it can be shown that they reduce neighbourhood problems related to consumption of drugs like crack cocaine and methamphetamine.

In contrast to Firestone-Cruz and colleagues [37], we created a composite measure of public opinion to highlight those with mixed opinions about SIFs. Focus group data from residents, business owners, and community service providers in our study also revealed high levels of ambivalence about SIF implementation [38]. We classified responses for each of the latter four questions into three categories: strongly disagree; somewhat agree or disagree (participants who might change their views); and strongly agree. Finally, we grouped responses to the four goals into a single composite variable using the following categories:

- Strongly agreed: participants who strongly agreed with all four goals for SSFs and SIFs
- Strongly disagreed: participants who strongly disagreed with all four goals for SSFs and SIFs
- Mixed opinions: all other patterns of responses

Analysis

We used statistical models for complex survey data, which account for stratified sampling and incorporate sampling weights. These weights reflect study design and population characteristics such that the final survey results are representative of the Ontario population aged 18 years in the survey year. Sampling weights also provide accurate confidence intervals. A full description of the analytic methods can be found in a previously published study [36]. We compared the sample who answered questions to the entire population sample and to data from the 2006 Ontario Census. We performed chi-square tests of independence, taking into account the complex survey design, to determine the association between SSF awareness and each of the four specific SSF goals. We used a two-tailed p-value threshold of 0.05 to assess statistical significance and did not adjust for multiple comparisons.

We analysed independent predictors of knowledge of SSF using logistic regression and of support for SSF goals using multinomial logistic regression, with a 3-

level dependent variable based on the composite support variable. Both methods used weights for complex survey designs. We used a non-parsimonious approach for each method and included all potential covariates listed in Table 1. Multinomial regression results are reported compared to the "mixed opinions" group as both the relative risk of strongly agreeing and of strongly disagreeing with the goals of SSFs.

All analyses were completed using SAS 9.3 (SAS Institute), R 3.0.2 (R Core Team), and Stata version 13.1 (StataCorp).

Results

In 2009, 1035 participants were asked questions about SIFs and SSFs. After applying survey weights, the typical respondent was female (55.0 %), aged 35 and over (72.2 %), employed full-time (62.4 %), and had completed at least some post-secondary education (68.1 %; Table 1). The participants who answered questions about SIFs and SSFs had a similar employment and income distribution as the Canadian population but women, older people, people who were married, and well-educated people were somewhat over-represented. At least one question regarding opinions about SIF or SSF implementation was not answered by 182 and 198 participants, respectively.

Using complex survey designs to estimate proportions, significantly fewer participants had ever read, seen or heard any information about SSFs (17.9 %, 95 % CI 15.1 % to 21.1 %) than SIFs (57.6 %, 95 % CI 53.9 % to 61.2 %); the difference was 39.7 % (95 % CI 35.8 % to 43.6 %; $p < 0.001$; difference estimated using the delta method using a t-distribution with 1010 degrees of freedom). Among those with no prior knowledge of SSFs, 48.8 % also had no prior knowledge of SIFs and among those with no prior knowledge of SIFs, 95.4 % also had no prior knowledge of SIFs thus 40.1 % of the population had no prior knowledge of either type of facility and 15.9 % had prior knowledge of both types (data not shown). Independent predictors of knowledge of SSFs included older age, male sex, having a university degree, and being a health care worker; people who immigrated to Canada were less likely have prior knowledge of SSFs (Table 2).

Participants had varied opinions about implementation of SSFs and SIFs. When asked if SSFs should be made available to people who smoke drugs such as crack cocaine and crystal methamphetamine, 64.2 % of Ontarians held a mixed overall opinion, 19.6 % strongly agreed, and 16.1 % strongly disagreed (Table 3). The distribution of responses was significantly different when participants were asked about SIFs; 28.3 % of participants strongly agreed and 11.6 % strongly disagreed. Analyses by each program goal showed significantly fewer participants

Table 1 Demographic characteristics^a

Characteristic	Weighted percent	Total survey	Census	Sample size; Design degrees of freedom	<i>p</i> -value; Chi-squared degrees of freedom
Female	55.0 %	51.5 %	51.9 %	1035; 1029	0.049; 1
Age (years)					
18 to 34	25.5 %	26.3 %	28.5 %	999; 993	0.012; 2
35 to 54	39.2 %	41.7 %	40.4 %		
55 and older	35.3 %	32.0 %	31.1 %		
Employment status					
Employed	63.2 %	63.7 %	64.5 %	1020; 1014	0.654; 2
Unemployed	4.5 %	5.1 %	4.1 %		
Other (student, retired, homemaker, disability, etc.)	32.3 %	31.1 %	31.4 %		
Household income in the past year before taxes					
Less than \$30,000	13.0 %	11.4 %	12.6 %	730; 724	0.952; 3
Between \$30,000 and \$49,999,99	16.1 %	15.6 %	16.6 %		
Between \$50,000 and \$79,999,99	24.1 %	26.0 %	24.6 %		
More than \$80,000	46.8 %	47.0 %	46.1 %		
Highest level of education attained					
Less than high school	10.5 %	10.6 %	18.6 %	1018; 1012	<0.001; 3
Completed high school	20.4 %	21.2 %	27.8 %		
Some post-secondary (college or university)	34.3 %	35.6 %	32.1 %		
University degree	34.8 %	32.7 %	21.4 %		
Marital Status					
Married/living with partner	71.0 %	69.1 %	62.9 %	1014; 1018	<0.001; 2
Previously married (divorced, widowed, separated)	10.6 %	10.6 %	14.1 %		
Never married	18.4 %	20.3 %	23.1 %		
Smoking Status			NA	1028; 1022	
Current	17.9 %	18.6 %			
Former	25.7 %	26.5 %			
Never	56.4 %	54.9 %			
Religious Service Attendance, past 12 months			NA	969; 963	
None	35.8 %	38.1 %			
1 to 6 times	29.3 %	27.7 %			
7 or more times	34.9 %	34.2 %			
Alcohol use in the past 12 months	79.0 %	79.1 %	NA	1031; 1025	
Cannabis use in the past 12 months	11.8 %	13.2 %	NA	1033; 1027	
Fair or Poor Self-reported Health	9.6 %	10.5 %	NA	1029; 1023	
Fair or Poor Self-reported Mental Health	6.2 %	5.7 %	NA	1025; 1019	
Health care worker	6.8 %	6.1 %	NA	908; 902	
Immigrant to Canada	27.2 %	27.5 %	NA	1014; 1008	
Urban residence	80.3 %	82.9 %	NA	1035; 1029	

^aProportions estimated from one-way tables using complex survey designs. Total survey sample size = 2037. Census estimates are from the 2006 Canadian Census individual microuse data (<http://www5.statcan.gc.ca/olc-cel/olc.action?ObjId=95M0028X&ObjType=2>) for Ontario population aged 18 and older using individual weighting factors. *P*-values compare values in the current sample to population estimates using Pearson's chi-square test. NA denotes not available

strongly agreed with implementation of SSFs than with implementation of SIFs (Table 3). For example, 20.5 % strongly agreed with implementation of SSFs to

encourage safer drug use compared to 30.6 % who strongly agreed with implementation of SIFs for that purpose. For each goal, significantly more participants

Table 2 Multivariable analysis of knowledge of supervised smoking facilities

Variable	Odds ratio (95 % Confidence Interval)
Age (per decade)	1.17 (1.02 to 1.34)
Male sex	1.88 (1.22 to 2.92)
Household income in the past year before taxes	
Less than \$30,000	1.00 (Referent)
Between \$30,000 and \$49,999,99	0.63 (0.31 to 1.29)
Between \$50,000 and \$79,999,99	0.57 (0.28 to 1.16)
More than \$80,000	0.91 (0.44 to 1.89)
Highest level of education attained	
Less than high school	1.00 (Referent)
Completed high school	0.75 (0.36 to 1.56)
Some post-secondary (college or	1.74 (0.85 to 3.58)
University degree	2.59 (1.16 to 5.78)
Marital Status	
Married/Living with partner	1.00 (Referent)
Previously married	0.84 (0.45 to 1.56)
Never married	1.33 (0.71 to 2.46)
Smoking Status	
Current	1.00 (Referent)
Former	1.04 (0.51 to 2.09)
Never	0.55 (0.31 to 0.99)
Religious Service Attendance, past 12 months	
None	1.00 (Referent)
1 to 6 times	0.62 (0.36 to 1.06)
7 or more times	0.88 (0.52 to 1.49)
Employment status	
Employed	1.00 (Referent)
Unemployed	1.31 (0.45 to 3.80)
Other (student, retired, homemaker)	1.18 (0.68 to 2.05)
Alcohol use in the past 12 months	0.84 (0.50 to 1.40)
Cannabis use in the past 12 months	1.25 (0.57 to 2.73)
Fair or Poor Self-reported Health	1.24 (0.62 to 2.51)
Fair or Poor Self-reported Mental Health	0.44 (0.17 to 1.10)
Health care worker	2.55 (1.08 to 6.01)
Immigrant to Canada	0.47 (0.28 to 0.81)
Urban residence	0.89 (0.56 to 1.44)

Odds ratios estimated using logistic regression for complex survey designs. Statistical significance assessed using t-distribution with 610 degrees of freedom

strongly disagreed with implementation of SSFs than strongly disagreed with implementation of SIFs.

Bivariate analyses showed that just over half of Ontarians (51.1 %) somewhat agreed/disagreed, 15.7 % strongly agreed and 10.6 % strongly disagreed with implementing both SSFs and SIFs (Table 4).

Prior knowledge of SSFs influenced support for implementation. Participants who had prior knowledge of SSFs were more likely to strongly agree with implementation of

SSFs on the composite measure of agreement than those who did not have prior knowledge (28.4 % vs. 16.8 %; difference 11.6 % [95 % CI 2.7 % to 20.5 %]) and less likely to report mixed opinions (55.1 % vs. 66.7 %; difference -11.6 % [95 % CI -22.1 % to -1.1 %]; Table 5). Participants with prior knowledge of SSFs were also more supportive of SSF implementation in relation to the goals of encouraging safer drug use (difference 12.8 % reducing infectious diseases (difference 14.9 % and slightly more likely to

Table 3 Difference in opinions about SSFs and SIFs by composite measure and specific goals

Goal of facility ^a	Supervised smoking	Supervised injection	Difference	p-value	Design Degrees of Freedom
	Weighted percent (95 % Confidence Interval)	Weighted percent (95 % Confidence Interval)	Weighted percent (95 % Confidence Interval)		
Composite measure					
Strongly agree	19.6 % (16.5 % to 23.2 %)	28.3 % (24.6 % to 32.3 %)	-8.7 % (-12.2 % to -5.2 %)	<0.001	772
Somewhat agree/disagree	64.2 % (60.1 % to 68.2 %)	60.1 % (55.9 % to 64.2 %)	4.1 % (0.1 % to 8.1 %)	0.046	
Strongly disagree	16.1 % (13.3 % to 19.4 %)	11.6 % (9.1 % to 14.6 %)	4.6 % (2.6 % to 6.5 %)	<0.001	
Supervised smoking facilities should be made available to people who smoke drugs like crack cocaine and methamphetamine to encourage safer drug use					
Strongly agree	20.5 % (17.3 % to 24%)	30.6 % (27.7 % to 34.5 %)	-10.1 % (-13.7 % to -6.6 %)	<0.001	838
Somewhat agree/disagree	40.5 % (36.5 % to 44.6 %)	40.3 % (36.3 % to 44.4 %)	0.2 % (-4.0 % to 4.3 %)	0.932	
Strongly disagree	39.1 % (35.2 % to 43.1 %)	29.1 % (25.6 % to 32.9 %)	10% (6.9 % to 13%)	<0.001	
Supervised smoking facilities should be made available if it can be shown that they reduce infectious disease among people who smoke drugs like crack cocaine and methamphetamine					
Strongly agree	35.2 % (31.4 % to 39.2 %)	49.4 % (45.4 % to 53.4 %)	-14.2 % (-17.7 % to -10.7 %)	<0.001	881
Somewhat agree/disagree	39.9 % (36.0 % to 43.9 %)	32.4 % (28.9 % to 36.2 %)	7.4 % (3.5 % to 11.4 %)	<0.001	
Strongly disagree	24.9 % (21.7 % to 28.4 %)	18.1 % (15.3 % to 21.4 %)	6.8 % (4.3 % to 9.3 %)	<0.001	
Supervised smoking facilities should be made available if they can increase drug users' contact with health and social workers					
Strongly agree	40.1 % (36.3 % to 44.1 %)	48.3 % (44.4 % to 52.3 %)	-8.2 % (-11.3 % to -5%)	<0.001	886
Somewhat agree/disagree	40% (36.2 % to 44%)	37.2 % (33.4 % to 41.1 %)	2.9 % (-0.5 % to 6.3 %)	0.097	
Strongly disagree	19.8 % (16.9 % to 23.1 %)	14.5 % (11.9 % to 17.5 %)	5.3 % (3.1 % to 7.5 %)	<0.001	
Supervised smoking facilities should be made available if it can be shown that they reduce neighbourhood problems related to use of drugs like crack cocaine and methamphetamine					
Strongly agree	45.7 % (41.8 % to 49.7 %)	56.1 % (52.2 % to 60%)	-10.4 % (-13.4 % to -7.4 %)	<0.001	910
Somewhat agree/disagree	36.6 % (32.9 % to 40.4 %)	31% (27.5 % to 34.8 %)	5.6 % (2.4 % to 8.7 %)	<0.001	
Strongly disagree	17.7 % (15% to 20.8 %)	12.9 % (10.5 % to 15.7 %)	4.8 % (3.1 % to 6.6 %)	<0.001	

^aSupervised injection questions were worded correspondingly to ask about drug injection. Proportions estimated using complex survey designs. Differences calculated using t-distribution for differences with specified degrees of freedom. For our survey design, sample size = design degrees of freedom + number of strata (6)

support implementation for increasing contact with health and social workers (difference 11.1 %), but not significantly associated with agreement to implement SSFs to reduce neighbourhood problems (difference 6.7 %). Strong disagreement with implementation of SSFs overall was not influenced by prior awareness of SSFs except for the goal of encouraging safer drug use, where participants who were not aware of SSFs were more likely to strongly disagree with implementation (41.4 % vs. 29.6 %).

Few variables were predictive of either strongly agreeing or strongly disagreeing with the goals of SSFs

(Table 6). People with an annual income between \$50,000 and \$80,000 and those who attended religious services 7 or more times in the year prior to being surveyed were significantly less likely to agree with the goals of SSFs compared to those who somewhat agreed/disagreed, whereas people who smoked cannabis in the past 12 months were significantly less likely to disagree with the goals of SSFs compared to those who somewhat agreed/disagreed. Post-estimation adjustment predicted for strongly disagreeing with the goals of SIFs for people with no, 1 to 6, and 7 or more religious service attendances per year were 9.3 %, 9.5 %, and 13.1 %.

Table 4 Bivariate analysis of composite measures of agreement regarding supervised injection and smoking facilities^a

Supervised injection composite measure	Supervised smoking composite measure			p-value
	Strongly agree	Somewhat agree/disagree	Strongly disagree	
Strongly Agree	15.7 %	12.1 %	0.5 %	
Somewhat agree/disagree	3.9 %	51.1 %	5.1 %	
Strongly disagree	0.0 %	1.0 %	10.6 %	<0.001

^aProportions estimated using complex survey designs. Test statistic for distribution uses Pearson's chi-square test converted to an F-statistic for the survey design with the second-order correction of Rao and Scott with $F(3.90, 3011.27) = 123.7119$ [46]

Table 5 Agreement with goals of a supervised smoking facility by level of awareness^a

Goal of facility	Aware of supervised smoking facilities					
	No (95 % CI)	Yes (95 % CI)	Difference (95 % CI)	p-value (difference)	p-value (distribution)	Degrees of freedom
Composite measure						
Strongly agree	16.8 % (13.6 % to 20.6 %)	28.4 % (21.0 % to 37.3 %)	11.6 % (2.7 % to 20.5 %)	0.011		Difference: 829 Distribution: F(1.99, 1652.12) = 4.0000
Somewhat agree/disagree	66.7 % (62.3 % to 70.8 %)	55.1 % (45.4 % to 64.4 %)	-11.6 % (-22.1 % to -1.1 %)	0.030		
Strongly disagree	16.5 % (13.5 % to 20.0 %)	16.5 % (10.5 % to 25.0 %)	0.0 % (-7.9 % to 7.9 %)	0.995	0.019	
Supervised smoking facilities should be made available to people who smoke drugs like crack cocaine and methamphetamine to encourage safer drug use						
Strongly agree	17.3 % (14.2 % to 21.0 %)	30.2 % (22.7 % to 38.9 %)	12.8 % (4.0 % to 21.6 %)	0.005		Difference: 889 Distribution: F(1.97, 1752.21) = 5.3906
Somewhat agree/disagree	41.2 % (37.0 % to 45.6 %)	40.3 % (31.0 % to 50.3 %)	-0.9 % (-11.6 % to 9.8 %)	0.863		
Strongly disagree	41.4 % (37.2 % to 45.7 %)	29.6 % (22.0 % to 38.5 %)	-11.9 % (-21.2 % to -2.5 %)	0.013	0.005	
Supervised smoking facilities should be made available if it can be shown that they reduce infectious disease among people who smoke drugs like crack cocaine and methamphetamine						
Strongly agree	31.8 % (27.8 % to 36.0 %)	46.7 % (37.4 % to 56.3 %)	14.9 % (4.5 % to 25.3 %)	0.005		Difference: 910 Distribution: F(1.98, 1802.95) = 4.4720
Somewhat agree/disagree	42.3 % (38.1 % to 46.6 %)	30.5 % (22.0 % to 40.5 %)	-11.8 % (-22.1 % to -1.6 %)	0.023		
Strongly disagree	25.9 % (22.4 % to 29.7 %)	22.8 % (16.0 % to 31.4 %)	-3.1 % (-11.6 % to 5.4 %)	0.478	0.012	
Supervised smoking facilities should be made available if they can increase drug users' contact with health and social workers						
Strongly agree	37.7 % (33.6 % to 42.0 %)	48.8 % (39.3 % to 58.4 %)	11.1 % (0.6 % to 21.6 %)	0.039		Difference: 919 Distribution: F(1.98, 1823.00) = 2.2802
Somewhat agree/disagree	42.1 % (37.9 % to 46.3 %)	33.1 % (24.3 % to 43.3 %)	-9.0 % (-19.5 % to 1.5 %)	0.093		
Strongly disagree	20.2 % (17.1 % to 23.8 %)	18.1 % (12.0 % to 26.4 %)	-2.1 % (-10.0 % to 5.8 %)	0.599	0.103	
Supervised smoking facilities should be made available if it can be shown that they reduce neighbourhood problems related to use of drugs like crack cocaine and methamphetamine						
Strongly agree	44.0 % (39.8 % to 48.2 %)	50.7 % (41.1 % to 60.2 %)	6.7 % (-3.8 % to 17.2 %)	0.213		Difference: 934 Distribution: F(1.97, 1844.31) = 1.1145
Somewhat agree/disagree	38.3 % (34.3 % to 42.4 %)	30.7 % (22.0 % to 40.9 %)	-7.6 % (-18.0 % to 2.8 %)	0.151		
Strongly disagree	17.8 % (14.9 % to 21.1 %)	18.7 % (12.5 % to 26.9 %)	0.9 % (-6.9 % to 8.8 %)	0.821	0.328	

^aCI denotes confidence interval. Proportions estimated using complex survey designs. Differences calculated using t-distribution for differences with specified degrees of freedom. Test statistic for distribution uses Pearson's chi-square test converted to an F-statistic for the survey design with the second-order correction of Rao and Scott with specified degrees of freedom [46]

Discussion

Although there are reports of decline in crack cocaine use in both Canada and the United States [39, 40], its use remains a public health concern. However, as noted above, public health programming for people who smoke crack cocaine lags behind what is available for people who inject drugs despite evidence of serious, and sometimes fatal, harms associated with smoking these drugs [41, 42]. To our knowledge, this is the first study to report public opinion about SSFs.

Lack of innovative public health responses to reduce stimulant drug-related problems parallels the lack of awareness about SSFs in the public opinion data. Our results show that less than 20 % of Ontarians were aware of SSF models whereas almost 60 % were aware of SIF models. SSF awareness was associated with older age, higher education and male sex. These results suggest the need for education among broad societal groups regarding supervised smoking. Compared to smoking facilities, we generally found stronger support for and fewer negative sentiments

Table 6 Multivariable analysis of agreement with the goals of supervised smoking facilities

Variable	Relative risk ratio (95 % Confidence Interval)	
	Strongly agree vs. somewhat agree/disagree	Strongly disagree vs. somewhat agree/disagree
Age (per decade)	1.09 (0.93 to 1.27)	0.93 (0.71 to 1.23)
Male sex	1.18 (0.71 to 1.99)	1.36 (0.64 to 2.89)
Household income in the past year before taxes		
Less than \$30,000	1.00 (Referent)	1.00 (Referent)
Between \$30,000 and \$49,999,99	0.73 (0.34 to 1.58)	0.91 (0.29 to 2.86)
Between \$50,000 and \$79,999,99	0.35 (0.15 to 0.81)	0.51 (0.17 to 1.52)
More than \$80,000	0.85 (0.38 to 1.93)	1.32 (0.36 to 4.82)
Highest level of education attained		
Less than high school	1.00 (Referent)	1.00 (Referent)
Completed high school	1.14 (0.46 to 2.82)	0.64 (0.24 to 1.71)
Some post-secondary (college or	0.74 (0.31 to 1.81)	0.46 (0.18 to 1.22)
University degree	1.52 (0.58 to 3.98)	0.43 (0.13 to 1.40)
Marital Status		
Married/Living with partner	1.00 (Referent)	1.00 (Referent)
Previously married	0.80 (0.37 to 1.72)	1.10 (0.43 to 2.80)
Never married	1.86 (0.91 to 3.77)	1.61 (0.57 to 4.55)
Smoking Status		
Current	1.00 (Referent)	1.00 (Referent)
Former	0.79 (0.37 to 1.67)	1.04 (0.36 to 2.99)
Never	0.90 (0.48 to 1.71)	0.78 (0.31 to 1.99)
Religious Service Attendance, past 12 months		
None	1.00 (Referent)	1.00 (Referent)
1 to 6 times	0.61 (0.34 to 1.08)	0.85 (0.30 to 2.36)
7 or more times	0.24 (0.13 to 0.45)	0.89 (0.37 to 2.16)
Employment status		
Employed	1.00 (Referent)	1.00 (Referent)
Unemployed	0.86 (0.28 to 2.67)	0.86 (0.14 to 5.06)
Other (student, retired, homemaker)	0.97 (0.53 to 1.76)	0.69 (0.26 to 1.81)
Alcohol use in the past 12 months	0.71 (0.36 to 1.37)	0.78 (0.35 to 1.75)
Cannabis use in the past 12 months	0.81 (0.38 to 1.72)	0.16 (0.03 to 0.76)
Fair or Poor Self-reported Health	0.96 (0.39 to 2.37)	1.03 (0.30 to 3.50)
Fair or Poor Self-reported Mental Health	0.68 (0.26 to 1.77)	0.66 (0.12 to 3.67)
Health care worker	1.37 (0.59 to 3.17)	0.21 (0.02 to 1.79)
Immigrant to Canada	0.83 (0.44 to 1.55)	1.49 (0.63 to 3.57)
Urban residence	1.12 (0.62 to 2.02)	0.71 (0.34 to 1.51)

Relative risk ratios estimated using multinomial logistic regression for complex survey designs. Statistical significance assessed using t-distribution with 536 degrees of freedom

about injection facilities. Support for SSFs was strongest when the goals of a facility were presented as reducing neighbourhood problems related to drug use, increasing contact between people who use drugs and health and social workers, and reducing infectious diseases. Even for these goals, however, we observed about 8 to 14 % higher

support for injection facilities, rather than smoking facilities. We found few consistent predictors of opposition to SSF goals although regular attendance at religious services was significant. These findings might indicate that both individual values and scepticism about the effectiveness of SSFs, reflecting the weaker

evidence base for SSFs compared to SIFs, underlie opposition. Our results underscore the need for ongoing research, including demonstration projects.

The survey was prone to sampling biases, as people who do not speak English or French and residents without a landline were excluded from participating. Nevertheless, the response rate was reasonable (57 %) for a telephone survey [35, 43] and the sampling was systematic. Thus, these data are population-based – unlike other studies assessing public opinion about SIFs that use convenience samples – and therefore more likely to be representative of true opinion, although highly educated people were somewhat over-represented, as commonly found in telephone surveys. The survey represents the last available data (in 2009), but SIFs have remained a prominent policy discussion topic since then. While our results might underestimate awareness of SIFs, they likely are still reflective of awareness of SSFs, which have been proposed but not implemented in Canada and are discussed much less frequently.

Grund and colleagues [22] note that stimulant use presents unique challenges requiring innovative and multidisciplinary responses and that people who use stimulants ‘must enjoy the fundamental human right to health protection, as stipulated by Article 25 of the Universal Declaration of Human Rights’ (p. 212). SSFs represent an innovation in programming that have been implemented in some European cities [9]. Until recently, harm reduction advocates in Canada have despaired over introducing innovations such as these given the opposition of the former Conservative federal government to harm reduction [44]. Again, our data also show low levels of public support for SSFs. Importantly, public opinion is identified in recent legislation, Bill C-2, passed in Canada that identified stakeholder opinion as one of the inputs that would be considered in any application for new SIFs [http://www.huffingtonpost.ca/2015/06/22/critics-up-in-arms-over-f_n_7640154.html]. However, public health advocates are hopeful that another SIF recently approved by the new federal government signals improved chances for approval of other applications for SIFs. [<http://www.theglobeandmail.com/news/british-columbia/vancouver-facility-becomes-canadas-second-approved-supervised-injection-site/article28216557/>]. The newly elected Prime Minister Justin Trudeau is on record expressing support for SIFs [<http://www.straight.com/blogra/404631/justin-trudeau-tells-ubc-students-he-wants-supervised-injection-sites-across-canada>], reflecting the leadership identified as necessary for innovation in programming and policy [45].

Conclusions

The lack of public knowledge or support of SSFs and their potential positive impact on public health issues associated with drug smoking presents barriers to service providers and community advocates who are contemplating implementation of SSFs in their jurisdiction.

Recent federal government changes in Canada may provide the leadership environment necessary to ensure that innovative harm reduction programs such as SSFs are developed based on evidence and implemented.

Abbreviations

SIF: Supervised injection facility: a common type of supervised consumption facility where people can inject illicitly obtained drugs; SSF: Supervised smoking facility: type of supervised consumption facility where people can smoke illicitly obtained drugs.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CS conceived of the study, participated in study design, assisted with statistical analyses, and drafted the first version of the manuscript. NKR conducted statistical analyses and assisted with preparation of the manuscript. TMW participated in study coordination and helped revise the manuscript. GK participated in study coordination and helped revise the manuscript. AMB conceived of the study, participated in study design, conducted statistical analyses, and helped revise the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank the full Toronto and Ottawa Supervised Consumption Assessment (TOSCA) team including the following co-investigators and team members: Patricia O'Campo, Richard Glazier, and Naushaba Degani from the Centre for Research on Inner City Health, St. Michael's Hospital; Benedikt Fischer from Simon Fraser University and CAMH; Shaun Hopkins from Toronto Public Health; Susan Shepherd from the Toronto Drug Strategy Secretariat; Janine Luce from CAMH; Peggy Millson from the University of Toronto; Gregory Zaric from the University of Western Ontario; Christopher Smith from the University of Pennsylvania; Lynne Leonard from the University of Ottawa; and Margaret Brandeau and Eva Enns from Stanford University. We would also like to thank the CAMH Monitor team for collecting the survey data. Funding support for this project was provided by the Ontario HIV Treatment Network and the Canadian Institutes of Health Research.

Author details

¹Dalla Lana School of Public Health, University of Toronto, 155 College Street, Toronto, Canada. ²Musculoskeletal Health and Outcomes Research, Li Ka Shing Knowledge Institute, St. Michael's Hospital, 30 Bond Street, Toronto, Canada. ³Centre for Research on Innercity Health, Li Ka Shing Knowledge Institute, St. Michael's Hospital, 30 Bond Street, Toronto, Canada. ⁴Department of Medicine, University of Toronto, 1 King's College Circle, Toronto, Canada. ⁵Institute of Health Policy, Management, and Evaluation, University of Toronto, 155 College Street, Toronto, Canada. ⁶Division of General Internal Medicine, St. Michael's Hospital, 30 Bond Street, Toronto, Canada.

Received: 29 July 2015 Accepted: 3 February 2016

Published online: 09 February 2016

References

- Marsch LA. The efficacy of methadone maintenance interventions in reducing illicit opiate use, HIV risk behavior and criminality: a meta-analysis. *Addiction*. 1998;93:515–32.
- Wodak A, Cooney A. Effectiveness of sterile needle and syringe programmes. *Int J Drug Policy*. 2005;16:531–44.
- Wood E(1), Werb D, Kazatchkine M, Kerr T, Hankins C, Gorna R, et al. Vienna Declaration: a call for evidence-based drug policies. *Lancet*. 2010;376(9738):310–2. doi:10.1016/S0140-6736(10)60958-0.
- Dolan K, Kimber J, Fry C, Fitzgerald J, McDonald D, Trautmann F. Drug consumption facilities in Europe and the establishment of supervised injecting centres in Australia. *Drug Alcohol Rev*. 2000;19:337–46.
- Bayoumi AM, Strike C, Jairam J, Watson T, Enns E, Kolla G, et al. Report of the Toronto and Ottawa Supervised Consumption Assessment Study. Toronto: St. Michael's Hospital and the Dalla Lana School of Public Health, University of Toronto; 2012.

6. Hedrich D. European report on drug consumption rooms. Lisbon: European Monitoring; 2004.
7. Kimber J, Dolan K, Van Beek I, Hedrich D, Zurhold H. Drug consumption facilities: an update since 2000. *Drug Alcohol Rev.* 2003;22:227–33.
8. Hedrich D, Kerr T, Dubois-Arber F. Drug consumption facilities in Europe and beyond. In: Rhodes T, Hedrich D. *Harm reduction: evidence, impacts and challenges*. Lisbon, Portugal: European Monitoring Centre for Drugs and Drug Addiction; 2010. p. 305–31.
9. Harm Reduction International. The global state of harm reduction 2014: towards an integrated response. London: Harm Reduction International; 2014.
10. Kerr T, Tyndall M, Li K, Montaner J, Wood E. Safer injecting facility use and syringe sharing in IDUs. *Lancet.* 2005;366:316–8.
11. Kimber J, Dolan K, Wodak A. Survey of drug consumption rooms: service delivery and perceived public health and amenity impact. *Drug Alcohol Rev.* 2005;24:21–4.
12. Marshall BD, Milloy M, Wood E, Montaner JS, Kerr T. Reduction in overdose mortality after the opening of North America's first medically supervised safer injecting facility: a retrospective population-based study. *Lancet.* 2011; 377:1429–37.
13. Potier C, Lapr evote V, Dubois-Arber F, Cottencin O, Rolland B. Supervised injection services: what has been demonstrated? A systematic literature review. *Drug Alcohol Depend.* 2014;145:48–68.
14. Salmon AM, Thein H, Kimber J, Kaldor JM, Maher L. Five years on: What are the community perceptions of drug-related public amenity following the establishment of the Sydney medically supervised injecting centre? *Int J Drug Policy.* 2007;18:46–53.
15. Wood E, Kerr T, Small W, Li K, Marsh DC, Montaner JSG, et al. Changes in public order after the opening of a medically supervised safer injecting facility for illicit injection drug users. *CMAJ.* 2004;171:731–4.
16. Wood E, Tyndall MW, Zhang R, Montaner JSG, Kerr T. Rate of detoxification service use and its impact among a cohort of supervised injecting facility users. *Addiction.* 2007;102:916–9.
17. Gyarmathy VA, Neaigus A, Miller M, Friedman SR, Des Jarlais DC. Risk correlates of prevalent HIV, hepatitis B virus, and hepatitis C virus infections among noninjecting heroin users. *J Acquir Immune Defic Syndr.* 2002;30:448–56.
18. Haydon E, Fischer B. Crack use as a public health problem in Canada: call for an evaluation of 'safer crack use kits'. *Can J Public Health.* 2005;96:185–8.
19. Leonard L, DeRubeis E, Pelude L, Medd E, Birkett N, Seto J. "I inject less as I have easier access to pipes". Injecting, and sharing of crack-smoking materials, decline as safer crack-smoking resources are distributed. *Int J Drug Policy.* 2008;19:255–64.
20. Porter J, Bonilla L. Crack users' cracked lips: an additional HIV risk factor. *Am J Public Health.* 1993;83:1490–1.
21. Tortu S, McMahon JM, Pouget ER, Hamid R. Sharing of noninjection drug-use implements as a risk factor for hepatitis C. *Subst Use Misuse.* 2004;39:211–.
22. Grund JP, Coffin P, Jauffret-Roustide M, Dijkstra M, de Bruin D, Blanken P. The fast and furious – cocaine, amphetamines and harm reduction. In: Rhodes T, Hedrich D, editors. *Harm reduction: evidence, impacts and challenges*. European Monitoring Centre for Drugs and Drug Addiction; 2010. p. 191–232.
23. Fischer B, Powis J, Cruz MF, Rudzinski K, Rehm J. Hepatitis C virus transmission among oral crack users: viral detection on crack paraphernalia. *Eur J Gastroen Hepat.* 2008;20:29–32.
24. Hermida M, Ferreira MC, Barral S, Laredo R, Castro A, Diz DP. Detection of HCV RNA in saliva of patients with hepatitis C virus infection by using a highly sensitive test. *J Virol Methods.* 2002;101:29–35.
25. McElroy PD, Rothenberg RB, Varghese R, Woodruff R, Minns GO, Muth SQ, et al. A network-informed approach to investigating a tuberculosis outbreak: implications for enhancing contact investigations. *Int J Tuberc Lung Dis.* 2003;7:5486–93.
26. McMahon JM, Simm M, Milano D, Clatts M. Detection of hepatitis C virus in the nasal secretions of an intranasal drug-user. *Ann Clin Microbiol Antimicrob.* 2004;3:6.
27. Strike C, Hopkins S, Watson TM, Gohil H, Leece P, Young S, et al. Best practice recommendations for Canadian harm reduction programs that provide service to people who use drugs and are at risk for HIV, HCV, and other harms: Part 1. Toronto: Working Group on Best Practice for Harm Reduction Programs in Canada; 2013.
28. Pizzey R, Hunt N. Distributing foil from needle and syringe programmes (NSPs) to promote transitions from heroin injecting to chasing: an evaluation. *Harm Reduct J.* 2008;524:1–8.
29. Collins CLC, Kerr T, Tyndall MW, Marsh DC, Kretz PS, Montaner JS, et al. Rationale to evaluate medically supervised safer smoking facilities for non-injection illicit drug users. *Can J Public Health.* 2005;96:344–7.
30. Shannon K, Ishida T, Morgan R, Bear A, Oleson M, Kerr T, et al. Potential community and public health impacts of medically supervised safer smoking facilities for crack cocaine users. *Harm Reduct J.* 2006;3.
31. McNeil R, Kerr T, Lampkin H, Small W. "We need somewhere to smoke crack: an ethnographic study of an unsanctioned safer smoking room in Vancouver, Canada. *Int J Drug Policy.* 2015;26:645–52.
32. Watson TM, Strike C, Kolla G, Penn R, Jairam J, Hopkins S, et al. Design considerations for supervised consumption facilities (SCFs): preferences for facilities where people can inject and smoke drugs. *Int J Drug Policy.* 2012; 24:156–63.
33. Burstein P. The impact of public opinion on public policy: a review and an agenda. *Polit Res Quart.* 2003;56:29–40.
34. Hyshka E, Bubela T, Wild TC. Prospects for scaling-up supervised injection facilities in Canada: the role of evidence in legal and political decision-making. *Addiction.* 2013;108:468–76.
35. Ialomiteanu A, Adlaf EM. CAMH Monitor 2009: technical guide. Toronto: Centre for Addiction and Mental Health; 2010.
36. Strike C, Jairam JA, Kolla G, Millson P, Shepherd S, Fischer B, et al. Increasing public support for supervised injection facilities in Ontario, Canada. *Addiction.* 2014;109:946–53.
37. Firestone-Cruz MF, Patra J, Fischer B, Rehm J, Kalousek K. Public opinion towards supervised injection facilities and heroin-assisted treatment in Ontario, Canada. *Int J Drug Policy.* 2007;18:54–61.
38. Strike C, Watson TM, Kolla G, Penn R, Bayoumi AM. Ambivalence about supervised injection facilities among community stakeholders. *Harm Reduct J.* 2015;12:26. doi:10.1186/s12954-015-0060-3.
39. Health Canada. Canadian alcohol and drug use monitoring survey: Summary of results for 2011. 2011. http://www.hc-sc.gc.ca/hc-ps/drugs-droguies/stat/_2011/summary-sommaire-eng.php. Accessed 6 Feb 2015.
40. Parker MA, Anthony JC. Should anyone be riding to glory on the now-descending limb of the crack-cocaine epidemic curve in the United States? *Drug Alcohol Depend.* 2014;138:225–8.
41. DeBeck K, Kerr T, Li K, Fischer B, Buxton J, Montaner J, et al. Smoking of crack cocaine as a risk factor for HIV infection among people who use injection drugs. *CMAJ.* 2009;181:585–9.
42. Leece P, Rajaram N, Woolhouse S, Millson M. Acute and chronic respiratory symptoms among primary care patients who smoke crack cocaine. *J Urban Health.* 2013;90:542–51.
43. Ialomiteanu A, Adlaf EM. CAMH Monitor 2003: technical guide. Toronto: Centre for Addiction and Mental Health; 2004.
44. Boyd N. Lessons from INSITE, Vancouver's supervised injection facility: 2003–2012. *Drugs Educ Prev Polic.* 2013;20:234–40.
45. Jauffret-Roustide M, Pedrono G, Beltzer N. Supervised consumption rooms: the French Paradox. *Int J Drug Policy.* 2013;24(6):628–30.
46. Rao JNK, Scott AJ. On chi-squared tests for multiway contingency tables with cell proportions estimated from survey data. *Ann Stat.* 1984;12:46–60.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

